

FECPANOVA, Yu. M. Cand Geolog-Mineralog Sci

Dissertation: "Morphology and Stratigraphy of the Cheilostomata
Neogene System of the Ponto-Caspian Basin." Moscow Order of the
Labor Red Banner Petroleum Inst imeni Academician I. M. Gubkin
10 Jun 47

SO: Vechernyaya Moskva, Jun 1947 (Proj #17836)

TEOFANOVA, Yu.M.

Study of upper Tertiary bryozoa of Moldavia and Crimea. *Biul.MOIP. Otd.geol.*
28 no.3:42-63 '53. (MLRA 6:11)

(Moldavia--Polysoa, Fossil) (Polysoa, Fossil--Moldavia)

(Crimea--Polysoa, Fossil) (Polysoa, Fossil--Crimea)

FEOFANOVA, Yu.M.

New Paleogene Bryozoa of Fergana. Paleont. zhur. no.1:60-72
165. (MIRA 18:4)

1. Moskovskiy institut neftekhimicheskoy i gazovoy promyshlennosti
imeni Gubkina.

1ST AND 2ND ORDERS										2ND AND 4TH ORDERS									
A B C D E F G H I J K L M N O P Q R S T U V W X Y Z										A B C D E F G H I J K L M N O P Q R S T U V W X Y Z									
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20										1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20									

PROCESS AND PROPERTIES INDEX

ECOFILDY, E.E.

Catalyst for the desulfurization of petroleum distillates.
E. E. Ecofilov, Russ. 43,702, July 31, 1935. An aq. soln. of $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$ is acidified with H_2SO_4 and SO_2 is passed through it at room temp. $\text{Cr}(\text{OH})_3$ is pptd. by addn. of NH_4OH and heated at 450° to produce Cr_2O_3 .

1ST AND 2ND ORDERS										2ND AND 4TH ORDERS									
A B C D E F G H I J K L M N O P Q R S T U V W X Y Z										A B C D E F G H I J K L M N O P Q R S T U V W X Y Z									
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20										1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20									

11 AND 120 ORDERS										11 AND 120 ORDERS									
PROCESSES AND PROPERTIES INDEX																			
FE OF FILDV E-1										18									
<p>Hydrogenation catalyst. V. Polozov and E. Feofilov. <i>Novosti Tekhniki</i> 1936, No. 36, 20.—Addn. of 20-5% of a portland cement to Mo sulfide catalyst increases its mech. stability. An aq. paste of this mixt. may be used for the prepn. of catalytic masses of various shapes. The use of an Fe frame for the paste increases the transfer and uniform distribution of heat. The method of the prepn. of Mo sul- fide from Na₂S and Mo oxides by treatment with water for 24 hr. and acidifying with dil. H₂SO₄ is worked out. A. A. Podgorny</p>																			
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>																			
11 AND 120 ORDERS										11 AND 120 ORDERS									

FEOFILOV, E.E.

250. CRITICAL OPERATIVE DE PROBLEMS OF THE POLICE, PLANNING, EXECUTION

FEOFILOV, G.L.

Some notes on the nomenclature of bronchi and bronchopulmonary segments. Arkh. anat., gist. i embr. 48 no.5: 102-104 My '65.

(MIRA 19:1)

1. Legochnoye otdeleniye (ispolnyayushchiy obyazannosti zav. - kand. med. nauk G.L. Feofilov) Instituta eksperimental'noy biologii i meditsiny Sibirskogo otdeleniya AN SSSR, Novosibirsk. Submitted January 31, 1963.

L 11933-66 EWT(1)/EWT(m)/T/EWP(t)/EWP(b)/EWA(c) IJP(c) JD/JG
ACC NR: AP6001654 SOURCE CODE: UR/0051/65/019/006/0973/0975

44 55 44 55 44 55
AUTHOR: Zonn, Z. N.; Ioffe, V. A.; Feofilov, P. O.

ORG: none

TITLE: Luminescence of chromium and manganese ions in lanthanum aluminate crystals

SOURCE: Optika i spektroskopiya, v. 19, no. 6, 1965, 973-975

TOPIC TAGS: manganese, chromium, ion, lanthanum compound, single crystal, luminescence, aluminate

ABSTRACT: The authors discuss certain results of the study of the spectra and luminescence duration of isoelectronic ions Cr^{3+} and Mn^{4+} (electronic configuration $3d^3$), introduced into the crystal lattice of LaAlO_3 . Both monocrystals grown from a solution in a melt as well as powdered samples were considered. No difference in the spectroscopic characteristics of the monocrystals and powders was noted. Crystal luminescence, located in the red and near-infrared portions of the spectrum was excited by an SVDSH-250 mercury lamp through a light filter consisting of a CuSO_4 solution, which had the effect of blocking the longwave portion of the energizing light. At small chromium concentrations

Card 1/2

UDC: 535.373.1

L 11933-66

ACC NR: AP6001654

APPROVED FOR RELEASE: 08/23/2000 CIA-RDP86-00513R000412910009-5"

the luminescence spectrum consists practically of a doublet with wavelengths of 7335 and 7338 Å. When the Cr^{3+} concentration is increased to 0.5% and above, the form of the luminescence spectrum undergoes definite modifications, which are described. With concentrations amounting to several percent, luminescence of Cr^{3+} in LaAlO_3 is entirely quenched. The luminescence spectra at different chromium percentages are illustrated. The luminescence is interpreted as a $3E - 4A_2$ transition in the Mn^{4+} ion. Orig. art. has: 3 figures.

SUB CODE: 20 / SUBM DATE: 20Mar65 / ORIG REF: 003 / OTH REF: 008

Card 2/2

15

FEDFAROVA, I. I.

CA

Determining the water-soluble minerals of saline soils. I. I. Fedfarova. *Pedology* (U. S. S. R.) 1940, No. 12, 40-52 (in German; 52). — With the aid of heavy liquids the minerals of the salinized soils have been sepd. The different sp. gr. fractions were examd. under the polarizing microscope and the minerals thus identified. In the Golodnaya steppe region the salt accumulations in the hard pan consist of halite and hemihydrate ($\text{CaSO}_4 \cdot 0.5\text{H}_2\text{O}$). Under the hard pan the hemihydrate disappears and gypsum is found. The white mellow hard pan contains primarily thenardite (Na_2SO_4). J. S. Joffe

ASIA-SLA METALLURGICAL LITERATURE CLASSIFICATION

CA FEOFAROVA, I. I.

Pseudomorphosis of calcite from gypsum in soils. I. I.
Feofarova. *Trudy Pochvennogo Inst. im. V. I. Dokuchaeva*
34, 302-6 (1950).--Field observations in the Fergan Valley
(Transcaucasia) and lab. expts. to test the field observa-
tions are reported. M. Hosen

CA FEOFAROVA, I. I.

Aragonite in soils. I. I. Feofarova. *Trudy Pochvennogo*
- *Izst. im. V. I. Dokuchaeva* 34, 20, 6 (1950). - Finding of
aragonite in certain soils of Transcaucasia is reported.
M. Hineh

1.07 0.0000, 1.1

Some data on the mineral composition of siliceous sprinkling in forest steppe soils. A. A. Roda and I. I. Feofarova. *Pochvovedenie* 1955, No. 9, 58-60. — The siliceous specks of a forest steppe soil were sepd. from the soil mass by scraping them with a sharp rod. Analysis of this material in comparison with that of the soil shows a concn. of SiO_2 in the specks or siliceous sprinkling, varying, resp., from 49 to 67% in the specks and from 33 to 43% in the soil. The feldspars (primarily orthoclase with some plagioclase) vary in the soil from 7 to 11% and in the specks from 11 to 14%. It is suggested that the mass of the specks represents the same material as that of the soil. However, the specks have lost their humus materials and clay aggregates. I. S. Joffe.

AG ①

Feofarova, I. I.

USSR/Soil Science - Physical and Chemical Properties of Soils.

J-3

Abs Jour : Ref Zhur - Biol., No 3, 1958, 10491

Author : Feofarova, I.I.

Inst :

Title : A Micromorphological Characterization of the Takyry

Orig Pub : Takyry Zap. Turkmenii i puti ikh s.-kh. osvoyeniya, Moskva, Akad Nauk SSSR, 1956, 351-380

Abstract : A study of the structure of slides taken from various horizons of the takyry of the Kopet-Dagh mountain plain, using a polarized microscope, has disclosed the cellular character of takyr porosity, the presence of gelatine-like newly-formed "secondary clay" in the lower part of the takyr crust, and also the existence of /osolodeniye/ spots in its upper part. Finely dispersed forms of carbonates were shown to coat the walls of the closed pores; the presence of some decarbonatized spots was also discovered. The great mobility of organic substances in the takyry is noted.

Card 1/1

FEOFAROVA, I.I.

~~Sulfates in saline soils. Trudy pochv.inst. 52:89-103 '58.~~

(MIRA 11:9)

(Alkali lands) (Sulfates)

TEOFAROVA, I.I.

Applying the microscopic method for determining carbonates in
saline soils. Trudy pochv.inst. 53:75-88 '58. (MIRA 11:9)
(Alkali lands) (Carbonates)

ZOMN, S.V.; PEOFAROVA, I.I.

New soil type under dark green conifer forests of eastern Tibet.
Pochvovedenie no.6:11-20 Je '60. (MIRA 13:11)

1. Laboratoriya lesovedeniya Akademii nauk SSSR.
(Tibet--Forest soils)

Concentration depolarization of the fluorescence of dye-
 stuff solutions. P. F. Fedorov and B. Ya. Sveshnikov.
J. Exptl. Theoret. Phys. (U. S. S. R.) 10, 1372-83 (1940);
J. Physics (U. S. S. R.) 3, 493-503 (1940). Exptl. data
 for fluorescein, eosin, rhodamin B, and trypanavin in
 glycerol and aq. solns. are shown in 18 figures. The de-
 polarization as found is represented by the Perrin equation
 (C. A. 24, 2673²) for rotational polarization with an addnl.
 term + Δc (c = concn.). P. H. Rathmann

[illegible]

FEOFILOV, P. P.

Inst. for Theor. Geoph., Acad. of Sci. USSR, Polucheno, (-1942-)

"Night Illumination and Spectral Composition of Night Light in Middle Latitudes,"

Iz. Ak. Nauk SSSR, Ser. Geograf. i Geofiz., Nos 1 - 6, 1942.

Theory of concentration depolarization of fluorescence in solutions. S. I. Vavilov and P. P. Kapitov. *Compt. rend. Acad. Sci. U.R.S.S.*, 1942, **84**, 220-223. — The relation $1/p' - 1/p = A/c$ (p' , p = degree of polarization of fluorescence respectively with and without concn. depolarization; τ = mean life of excited state; c = solute concn.; A const.) is deduced theoretically on the basis of excitation energy transfer by quantum-mechanical resonance.

A. J. E. W.

26

7.1.1940 - 1942 - 26

Polarization of fluorescence and anisotropy of molecules of organic dyes. P. P. Kuznetsov (J. Physics U.S.S.R., 1943, 7, 68-79).—Curves showing the dependence of polarization of fluorescent light on the λ of the exciting light (polarization spectra) are obtained for rhodamine-B, Na fluorescein, Na eosin (I), trypanflavine (II), acridine-orange (III), benzodiazine (IV), Na perylene-3,4,9,10-tetracarboxylate, mercuric orange (III), benzodiazine (IV). These polarization spectra are sp. for each dye. For (I)–(IV) the variation of dichroism of oriented mole. with λ is investigated; and there is a correspondence between this and the polarization spectrum. This indicates that the angles between emitting and absorbing oscillators are not produced during the excitation, but are inherent in the mol. The anisotropy of the mole. can be interpreted by means of an oscillator model in which the fluorescing mol. is regarded as an emitting oscillator with a no. of absorbing oscillators rigidly coupled to it.

A. J. M.

1ST AND 2ND ORDERS																										PROCESSES AND PROPERTIES INDEX																									
<p>Polarized luminescence of dyestuffs in rigid solutions. P. P. Fomilov. <i>Compt. rend. acad. sci. U.R.S.S.</i> 44, 147- 51; <i>Soviet Acad. Nauk S.S.S.R.</i> 44, 150-54(1944).— Measurements of polarization curves were carried out in parallel for the phosphorescence of a rigid soln. (I) and for the instantaneous fluorescence (II) of a glycerol-water soln. of the same dye. Both correspond to elec. di- pole radiation. In the majority of investigated cases the polarization spectra of I and II prove to be identical, giving evidence of the identity of the anisotropy of "centers" responsible for both luminescences. This is also addit. evidence that the characteristic dependence of polariza- tion on the wave length of the exciting light is not due to the oscillator's changing its orientation during the exist- ence of the excited state. The identity of concentrational curves for I and II proves that the processes which det. concentrational depolarization and quenching do not take place while the mol. is in the metastable state, or that they proceed with negligible probability. Thus, the energy transfer between an unexcited mol. and a mol. in the phos- phorescent state is impossible. Only after passing to the labile state is energy transfer possible. H. G. McCann</p>																																																			
<p>ASO-55 A METALLURGICAL LITERATURE CLASSIFICATION</p>																																																			
<p>GROUPS</p>																																																			

Fluorescence of organic dyes as related to chemical constitution.
P. P. Faglavay (*Compt. rend. Acad. Sci. U.R.S.S.,* 1946, 48, 367.
370).—Fluoregenic action is observed when dyes the structure of which admits of relative rotation of two large mol parts around the chain of conjugated double bonds responsible for the colour of the compound, are introduced into rigid solutions or sugar, plastic,
H₂O,, etc. Examples are di- and tri-phenylmethane dyes such as Auramine, fuchsin, crystal-violet, and malachite-green, and cyanine
dyes such as cyanine, dicyanine, and pinaverolinol. It is possible that some hindrance arises to preclude the relative motion of 1½ or more quinoline rings around the connecting chain of conjugated double bonds. This hindrance prevents the energy of excitation being converted into kinetic energy, and this view of fluorescence is preferred to the usual one in which fluorescence is ascribed as directly due to the presence of certain groupings.
C. R. H.

C. R. 11


ASME METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
PROCESSES AND PROPERTIES INDEX																																																			
<p><i>Fluorescence of organic dyes as related to chemical constitution.</i> P. P. Feofilov. <i>Doklady Akad. Nauk S.S.S.R.</i> 45, 387-90(1944); <i>Comp. rend. acad. sci. U.R.S.S.</i> 45, 367-70(1944)(in English).—Strong fluorescence may be imparted to nonfluorescent triphenylmethane (TPM) dyes either (1) by formation of a chem. bridge between 2 of the benzene rings, i.e., by conversion to acridine, xanthene, or similar dyes, or (2) by incorporating the TPM dyes in rigid solns., e.g., sugar, plastics, boric acid, or (3) by fixation on certain silicates. These methods of imparting fluorescence are effective with all dyes, e.g., auramine, diphenylmethane, TPM, cyanine, phneverdol, dicyanine, whose structure permits relative rotation of two large parts of the mol. about a chain of conjugated double bonds responsible for the color of the dye. It is suggested that the above-mentioned methods for imparting fluorescence are effective because they prevent aromatic groups in the dye mol. from undergoing an angular shift in position about the ethylene double bond in the excited state. Suppression of such rotation in effect seals energy within the mol. and thus makes the energy available for emission as fluorescent light. J. W. Perry.</p>																																																			
<p>ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>																																																			

SECHENOV, A. N. : FEOTILOV, P. P.

Luminescent materials accepted in a military sense.

An article found in "Optics for Military Use", Part I, published by the USSR Academy of Science, Moscow, 1945.



FEOFILOV, P.P.

PA 50198

USSR/Physics

Luminescence

Light - Polarization

Feb 1947

"Nature of Elementary Emitters and the Polarization of Photoluminescence," P. P. Feofilov, Lab Luminescence, State Opt Inst, 4 pp

"Camp Rend Acad Sci USSR" Vol IV, No 5

Vavilov has shown that the kind of multipolarity of elementary absorbing and emitting systems in the case of complicated luminescent molecules can be established by studying absorption and emission as function of the direction of the exciting light vector, and by

50198

USSR/Physics (Contd)

Feb 1947

observation. He has computed polarization diagrams for the four possible combinations of electric dipoles and quadrupoles and made it clear that the study of polarization diagrams permits definite decisions on the kind of multipolarity of elementary emitters. Feofilov extends Vavilov's considerations to the case of a magnetic dipole. Submitted by S. I. Vavilov, 19 Oct 1946.

50198

CA 3

Structure of the absorption band of triphenylmethane dyes. P. P. Perfilov. *Doklady Akad. Nauk S.S.S.R.* 57, 447-55 (1947).—The method of polarization spectra is used to indicate that in the dyes with a nearly sym. structure the secondary abs. max. are detd. by the structures of the mol. and are connected with the existence of an electronic shift which is different from that detg. the principal absorption band. Excitation of luminescence in the region of the long-wave absorption band gives the limiting polarization that is about 0.5 of the value calcd. for linear oscillators with coincident direction of oscillation, for all the previously studied dyes. Studies with fuchsin, pararufusin, methyl violet, crystal violet, ethyl violet, alk. phenolphthalein, and malachite green showed that dyes with 2 auxochromes display polarization that is close to the limiting value and is essentially invariant in respect to frequency of exciting light; here the principal oscillator system must be coaxial with the line joining the 2 auxochromes; these dyes do not show a secondary abs. maxima. Dyes with 3 auxochromes display a secondary abs. max. and the extent of polarization changes sharply with change of frequency of incident light within the limits of the absorption band; excitation with light of frequency corresponding to the long-wave edge of the band polarization is the same as cited above (40-50%), but with shorter waves the polarization falls off rapidly and may become neg. (abs. value 5-7%). G. M. Kosolapoff

PA 53T91

FEOFILOV, P. P.

USSR/Physics
Luminescent Materials
Polarization

Aug 1947

"Polarization of Photoluminescence and the Symmetry of the Molecule," P. P. Feofilov, Luminescence Lab, State Optical Inst, 4 pp

"Dok Akad Nauk SSSR, Nova Ser" Vol LVII, No 4

Presents in tabular form polarization of luminescence of benzole derivatives, important because of increased use of organic dyes. Analysis of molecule structure of fluorescent dyes shows that majority of complex molecules can be isolated and divided in several ways. Submitted by Academician S. I. Vavilov, 30 Mar 1947.

53T91

SA

533.37.08 - 52

2784

The study of the decay of luminescence by means of a
cathode ray oscilloscope. Tsvetkov, M. A. AND PROKOPEV,
P. R. *C.R. Acad. Sci. USSR*, 20-4/6, 3) 380-82 (1947) in
Russian. Translated into English. The phosphorescence with
cathode ray oscilloscope using exponential time-
base. The time constant of the timebase circuit is
constant (10⁻⁶ to 10⁻⁷ sec) of the timebase circuit is
adjustable and gives a linear trace when equal to the
decay time constant of the phosphor. The method can
be extended to non-exponential decay laws, for both
flash and afterglow. M. G. G.

PROCESSING AND PROPERTIES INDEX

1ST AND 2ND ORDERS

3RD AND 4TH ORDERS

5TH AND 6TH ORDERS

7TH AND 8TH ORDERS

9TH AND 10TH ORDERS

11TH AND 12TH ORDERS

13TH AND 14TH ORDERS

15TH AND 16TH ORDERS

17TH AND 18TH ORDERS

19TH AND 20TH ORDERS

21ST AND 22ND ORDERS

23RD AND 24TH ORDERS

25TH AND 26TH ORDERS

27TH AND 28TH ORDERS

29TH AND 30TH ORDERS

31ST AND 32ND ORDERS

33RD AND 34TH ORDERS

35TH AND 36TH ORDERS

37TH AND 38TH ORDERS

39TH AND 40TH ORDERS

41ST AND 42ND ORDERS

43RD AND 44TH ORDERS

45TH AND 46TH ORDERS

47TH AND 48TH ORDERS

49TH AND 50TH ORDERS

51ST AND 52ND ORDERS

53RD AND 54TH ORDERS

55TH AND 56TH ORDERS

57TH AND 58TH ORDERS

59TH AND 60TH ORDERS

61ST AND 62ND ORDERS

63RD AND 64TH ORDERS

65TH AND 66TH ORDERS

67TH AND 68TH ORDERS

69TH AND 70TH ORDERS

71ST AND 72ND ORDERS

73RD AND 74TH ORDERS

75TH AND 76TH ORDERS

77TH AND 78TH ORDERS

79TH AND 80TH ORDERS

81ST AND 82ND ORDERS

83RD AND 84TH ORDERS

85TH AND 86TH ORDERS

87TH AND 88TH ORDERS

89TH AND 90TH ORDERS

91ST AND 92ND ORDERS

93RD AND 94TH ORDERS

95TH AND 96TH ORDERS

97TH AND 98TH ORDERS

99TH AND 100TH ORDERS

101ST AND 102ND ORDERS

103RD AND 104TH ORDERS

105TH AND 106TH ORDERS

107TH AND 108TH ORDERS

109TH AND 110TH ORDERS

111ST AND 112ND ORDERS

113RD AND 114TH ORDERS

115TH AND 116TH ORDERS

117TH AND 118TH ORDERS

119TH AND 120TH ORDERS

121ST AND 122ND ORDERS

123RD AND 124TH ORDERS

125TH AND 126TH ORDERS

127TH AND 128TH ORDERS

129TH AND 130TH ORDERS

131ST AND 132ND ORDERS

133RD AND 134TH ORDERS

135TH AND 136TH ORDERS

137TH AND 138TH ORDERS

139TH AND 140TH ORDERS

141ST AND 142ND ORDERS

143RD AND 144TH ORDERS

145TH AND 146TH ORDERS

147TH AND 148TH ORDERS

149TH AND 150TH ORDERS

151ST AND 152ND ORDERS

153RD AND 154TH ORDERS

155TH AND 156TH ORDERS

157TH AND 158TH ORDERS

159TH AND 160TH ORDERS

161ST AND 162ND ORDERS

163RD AND 164TH ORDERS

165TH AND 166TH ORDERS

167TH AND 168TH ORDERS

169TH AND 170TH ORDERS

171ST AND 172ND ORDERS

173RD AND 174TH ORDERS

175TH AND 176TH ORDERS

177TH AND 178TH ORDERS

179TH AND 180TH ORDERS

181ST AND 182ND ORDERS

183RD AND 184TH ORDERS

185TH AND 186TH ORDERS

187TH AND 188TH ORDERS

189TH AND 190TH ORDERS

191ST AND 192ND ORDERS

193RD AND 194TH ORDERS

195TH AND 196TH ORDERS

197TH AND 198TH ORDERS

199TH AND 200TH ORDERS

201ST AND 202ND ORDERS

203RD AND 204TH ORDERS

205TH AND 206TH ORDERS

207TH AND 208TH ORDERS

209TH AND 210TH ORDERS

211ST AND 212ND ORDERS

213RD AND 214TH ORDERS

215TH AND 216TH ORDERS

217TH AND 218TH ORDERS

219TH AND 220TH ORDERS

221ST AND 222ND ORDERS

223RD AND 224TH ORDERS

225TH AND 226TH ORDERS

227TH AND 228TH ORDERS

229TH AND 230TH ORDERS

231ST AND 232ND ORDERS

233RD AND 234TH ORDERS

235TH AND 236TH ORDERS

237TH AND 238TH ORDERS

239TH AND 240TH ORDERS

241ST AND 242ND ORDERS

243RD AND 244TH ORDERS

245TH AND 246TH ORDERS

247TH AND 248TH ORDERS

249TH AND 250TH ORDERS

251ST AND 252ND ORDERS

253RD AND 254TH ORDERS

255TH AND 256TH ORDERS

257TH AND 258TH ORDERS

259TH AND 260TH ORDERS

261ST AND 262ND ORDERS

263RD AND 264TH ORDERS

265TH AND 266TH ORDERS

267TH AND 268TH ORDERS

269TH AND 270TH ORDERS

271ST AND 272ND ORDERS

273RD AND 274TH ORDERS

275TH AND 276TH ORDERS

277TH AND 278TH ORDERS

279TH AND 280TH ORDERS

281ST AND 282ND ORDERS

283RD AND 284TH ORDERS

285TH AND 286TH ORDERS

287TH AND 288TH ORDERS

289TH AND 290TH ORDERS

291ST AND 292ND ORDERS

293RD AND 294TH ORDERS

295TH AND 296TH ORDERS

297TH AND 298TH ORDERS

299TH AND 300TH ORDERS

301ST AND 302ND ORDERS

303RD AND 304TH ORDERS

305TH AND 306TH ORDERS

307TH AND 308TH ORDERS

309TH AND 310TH ORDERS

311ST AND 312ND ORDERS

313RD AND 314TH ORDERS

315TH AND 316TH ORDERS

317TH AND 318TH ORDERS

319TH AND 320TH ORDERS

321ST AND 322ND ORDERS

323RD AND 324TH ORDERS

325TH AND 326TH ORDERS

327TH AND 328TH ORDERS

329TH AND 330TH ORDERS

331ST AND 332ND ORDERS

333RD AND 334TH ORDERS

335TH AND 336TH ORDERS

337TH AND 338TH ORDERS

339TH AND 340TH ORDERS

341ST AND 342ND ORDERS

343RD AND 344TH ORDERS

345TH AND 346TH ORDERS

347TH AND 348TH ORDERS

349TH AND 350TH ORDERS

351ST AND 352ND ORDERS

353RD AND 354TH ORDERS

355TH AND 356TH ORDERS

357TH AND 358TH ORDERS

359TH AND 360TH ORDERS

361ST AND 362ND ORDERS

363RD AND 364TH ORDERS

365TH AND 366TH ORDERS

367TH AND 368TH ORDERS

369TH AND 370TH ORDERS

371ST AND 372ND ORDERS

373RD AND 374TH ORDERS

375TH AND 376TH ORDERS

377TH AND 378TH ORDERS

379TH AND 380TH ORDERS

381ST AND 382ND ORDERS

383RD AND 384TH ORDERS

385TH AND 386TH ORDERS

387TH AND 388TH ORDERS

389TH AND 390TH ORDERS

391ST AND 392ND ORDERS

393RD AND 394TH ORDERS

395TH AND 396TH ORDERS

397TH AND 398TH ORDERS

399TH AND 400TH ORDERS

401ST AND 402ND ORDERS

403RD AND 404TH ORDERS

405TH AND 406TH ORDERS

407TH AND 408TH ORDERS

409TH AND 410TH ORDERS

411ST AND 412ND ORDERS

413RD AND 414TH ORDERS

415TH AND 416TH ORDERS

417TH AND 418TH ORDERS

419TH AND 420TH ORDERS

421ST AND 422ND ORDERS

423RD AND 424TH ORDERS

425TH AND 426TH ORDERS

427TH AND 428TH ORDERS

429TH AND 430TH ORDERS

431ST AND 432ND ORDERS

433RD AND 434TH ORDERS

435TH AND 436TH ORDERS

437TH AND 438TH ORDERS

439TH AND 440TH ORDERS

441ST AND 442ND ORDERS

443RD AND 444TH ORDERS

445TH AND 446TH ORDERS

447TH AND 448TH ORDERS

449TH AND 450TH ORDERS

451ST AND 452ND ORDERS

453RD AND 454TH ORDERS

455TH AND 456TH ORDERS

457TH AND 458TH ORDERS

459TH AND 460TH ORDERS

461ST AND 462ND ORDERS

463RD AND 464TH ORDERS

465TH AND 466TH ORDERS

467TH AND 468TH ORDERS

469TH AND 470TH ORDERS

471ST AND 472ND ORDERS

473RD AND 474TH ORDERS

475TH AND 476TH ORDERS

477TH AND 478TH ORDERS

479TH AND 480TH ORDERS

481ST AND 482ND ORDERS

483RD AND 484TH ORDERS

485TH AND 486TH ORDERS

487TH AND 488TH ORDERS

489TH AND 490TH ORDERS

491ST AND 492ND ORDERS

493RD AND 494TH ORDERS

495TH AND 496TH ORDERS

497TH AND 498TH ORDERS

499TH AND 500TH ORDERS

501ST AND 502ND ORDERS

503RD AND 504TH ORDERS

505TH AND 506TH ORDERS

507TH AND 508TH ORDERS

509TH AND 510TH ORDERS

511ST AND 512ND ORDERS

513RD AND 514TH ORDERS

515TH AND 516TH ORDERS

517TH AND 518TH ORDERS

519TH AND 520TH ORDERS

521ST AND 522ND ORDERS

523RD AND 524TH ORDERS

525TH AND 526TH ORDERS

527TH AND 528TH ORDERS

529TH AND 530TH ORDERS

531ST AND 532ND ORDERS

533RD AND 534TH ORDERS

535TH AND 536TH ORDERS

537TH AND 538TH ORDERS

539TH AND 540TH ORDERS

541ST AND 542ND ORDERS

543RD AND 544TH ORDERS

545TH AND 546TH ORDERS

547TH AND 548TH ORDERS

549TH AND 550TH ORDERS

551ST AND 552ND ORDERS

553RD AND 554TH ORDERS

555TH AND 556TH ORDERS

557TH AND 558TH ORDERS

559TH AND 560TH ORDERS

561ST AND 562ND ORDERS

563RD AND 564TH ORDERS

565TH AND 566TH ORDERS

567TH AND 568TH ORDERS

569TH AND 570TH ORDERS

571ST AND 572ND ORDERS

573RD AND 574TH ORDERS

575TH AND 576TH ORDERS

577TH AND 578TH ORDERS

579TH AND 580TH ORDERS

581ST AND 582ND ORDERS

583RD AND 584TH ORDERS

585TH AND 586TH ORDERS

587TH AND 588TH ORDERS

589TH AND 590TH ORDERS

591ST AND 592ND ORDERS

593RD AND 594TH ORDERS

595TH AND 596TH ORDERS

597TH AND 598TH ORDERS

599TH AND 600TH ORDERS

601ST AND 602ND ORDERS

603RD AND 604TH ORDERS

605TH AND 606TH ORDERS

607TH AND 608TH ORDERS

609TH AND 610TH ORDERS

611ST AND 612ND ORDERS

613RD AND 614TH ORDERS

615TH AND 616TH ORDERS

617TH AND 618TH ORDERS

619TH AND 620TH ORDERS

621ST AND 622ND ORDERS

623RD AND 624TH ORDERS

625TH AND 626TH ORDERS

627TH AND 628TH ORDERS

629TH AND 630TH ORDERS

631ST AND 632ND ORDERS

633RD AND 634TH ORDERS

635TH AND 636TH ORDERS

637TH AND 638TH ORDERS

639TH AND 640TH ORDERS

641ST AND 642ND ORDERS

643RD AND 644TH ORDERS

645TH AND 646TH ORDERS

647TH AND 648TH ORDERS

649TH AND 650TH ORDERS

651ST AND 652ND ORDERS

653RD AND 654TH ORDERS

655TH AND 656TH ORDERS

657TH AND 658TH ORDERS

659TH AND 660TH ORDERS

661ST AND 662ND ORDERS

663RD AND 664TH ORDERS

665TH AND 666TH ORDERS

667TH AND 668TH ORDERS

669TH AND 670TH ORDERS

671ST AND 672ND ORDERS

673RD AND 674TH ORDERS

675TH AND 676TH ORDERS

677TH AND 678TH ORDERS

679TH AND 680TH ORDERS

681ST AND 682ND ORDERS

683RD AND 684TH ORDERS

685TH AND 686TH ORDERS

687TH AND 688TH ORDERS

689TH AND 690TH ORDERS

691ST AND 692ND ORDERS

693RD AND 694TH ORDERS

695TH AND 696TH ORDERS

697TH AND 698TH ORDERS

699TH AND 700TH ORDERS

701ST AND 702ND ORDERS

703RD AND 704TH ORDERS

705TH AND 706TH ORDERS

707TH AND 708TH ORDERS

709TH AND 710TH ORDERS

711ST AND 712ND ORDERS

713RD AND 714TH ORDERS

715TH AND 716TH ORDERS

717TH AND 718TH ORDERS

719TH AND 720TH ORDERS

721ST AND 722ND ORDERS

723RD AND 724TH ORDERS

725TH AND 726TH ORDERS

727TH AND 728TH ORDERS

729TH AND 730TH ORDERS

731ST AND 732ND ORDERS

733RD AND 734TH ORDERS

735TH AND 736TH ORDERS

737TH AND 738TH ORDERS

739TH AND 740TH ORDERS

741ST AND 742ND ORDERS

743RD AND 744TH ORDERS

745TH AND 746TH ORDERS

747TH AND 748TH ORDERS

749TH AND 750TH ORDERS

751ST AND 752ND ORDERS

753RD AND 754TH ORDERS

755TH AND 756TH ORDERS

757TH AND 758TH ORDERS

759TH AND 760TH ORDERS

761ST AND 762ND ORDERS

763RD AND 764TH ORDERS

765TH AND 766TH ORDERS

767TH AND 768TH ORDERS

769TH AND 770TH ORDERS

771ST AND 772ND ORDERS

773RD AND 774TH ORDERS

775TH AND 776TH ORDERS

777TH AND 778TH ORDERS

779TH AND 780TH ORDERS

781ST AND 782ND ORDERS

783RD AND 784TH ORDERS

785TH AND 786TH ORDERS

787TH AND 788TH ORDERS

789TH AND 790TH ORDERS

791ST AND 792ND ORDERS

793RD AND 794TH ORDERS

795TH AND 796TH ORDERS

797TH AND 798TH ORDERS

799TH AND 800TH ORDERS

801ST AND 802ND ORDERS

803RD AND 804TH ORDERS

805TH AND 806TH ORDERS

807TH AND 808TH ORDERS

809TH AND 810TH ORDERS

811ST AND 812ND ORDERS

813RD AND 814TH ORDERS

815TH AND 816TH ORDERS

817TH AND 818TH ORDERS

819TH AND 820TH ORDERS

821ST AND 822ND ORDERS

823RD AND 824TH ORDERS

825TH AND 826TH ORDERS

827TH AND 828TH ORDERS

829TH AND 830TH ORDERS

831ST AND 832ND ORDERS

833RD AND 834TH ORDERS

835TH AND 836TH ORDERS

837TH AND 838TH ORDERS

839TH AND 840TH ORDERS

841ST AND 842ND ORDERS

843RD AND 844TH ORDERS

845TH AND 846TH ORDERS

847TH AND 848TH ORDERS

849TH AND 850TH ORDERS

851ST AND 852ND ORDERS

853RD AND 854TH ORDERS

855TH AND 856TH ORDERS

857TH AND 858TH ORDERS

859TH AND 860TH ORDERS

861ST AND 862ND ORDERS

863RD AND 864TH ORDERS

865TH AND 866TH ORDERS

867TH AND 868TH ORDERS

869TH AND 870TH ORDERS

871ST AND 872ND ORDERS

873RD AND 874TH ORDERS

875TH AND 876TH ORDERS

877TH AND 878TH ORDERS

879TH AND 880TH ORDERS

881ST AND 882ND ORDERS

883RD AND 884TH ORDERS

885TH AND 886TH ORDERS

887TH AND 888TH ORDERS

889TH AND 890TH ORDERS

891ST AND 892ND ORDERS

893RD AND 894TH ORDERS

895TH AND 896TH ORDERS

897TH AND 898TH ORDERS

899TH AND 900TH ORDERS

901ST AND 902ND ORDERS

903RD AND 904TH ORDERS

905TH AND 906TH ORDERS

907TH AND 908TH ORDERS

909TH AND 910TH ORDERS

911ST AND 912ND ORDERS

913RD AND 914TH ORDERS

915TH AND 916TH ORDERS

917TH AND 918TH ORDERS

919TH AND 920TH ORDERS

921ST AND 922ND ORDERS

923RD AND 924TH ORDERS

925TH AND 926TH ORDERS

927TH AND 928TH ORDERS

929TH AND 930TH ORDERS

931ST AND 932ND ORDERS

933RD AND 934TH ORDERS

935TH AND 936TH ORDERS

937TH AND 938TH ORDERS

939TH AND 940TH ORDERS

941ST AND 942ND ORDERS

943RD AND 944TH ORDERS

945TH AND 946TH ORDERS

947TH AND 948TH ORDERS

949TH AND 950TH ORDERS

951ST AND 952ND ORDERS

953RD AND 954TH ORDERS

955TH AND 956TH ORDERS

957TH AND 958TH ORDERS

959TH AND 960TH ORDERS

961ST AND 962ND ORDERS

963RD AND 964TH ORDERS

965TH AND 966TH ORDERS

967TH AND 968TH ORDERS

969TH AND 970TH ORDERS

971ST AND 972ND ORDERS

973RD AND 974TH ORDERS

975TH AND 976TH ORDERS

977TH AND 978TH ORDERS

979TH AND 980TH ORDERS

981ST AND 982ND ORDERS

983RD AND 984TH ORDERS

985TH AND 986TH ORDERS

987TH AND 988TH ORDERS

989TH AND 990TH ORDERS

991ST AND 992ND ORDERS

993RD AND 994TH ORDERS

995TH AND 996TH ORDERS

997TH AND 998TH ORDERS

999TH AND 1000TH ORDERS

FEOFILOV, P. P.

Apr 48

USSR/Physics
Light - Measurements
Optics, Physical

"One Possibility of an Equation of Depth of Modulation in the Kerr Effect," N. A. Tolstoy,
P. P. Feofilov, State Opt Inst, 3 pp

"Dok Akad Nauk SSSR, Nova Ser" Vol LX, No 2

Describes fundamentals of method for measuring Kerr Effect. Presents method for
calculating intensity of light passing through the analyzer. Submitted by Academician
S. I. Vavilov, 18 Feb 1948.

PA 62197

FEOFILOV, P. P.

USSR/ Physics
Luminescence
Glass

11 Jan 1948

"Loss of Luminescence by Glass and Uranyl Salts" N. A. Tolstoy, P. P.
Feofilov, Luminescence Lab, State Opt Inst, 4 pp

"Dok Akad Nauk SSSR, Nova Ser" Vol LIX, No 2

Presents some results of use of new method to study attenuation of luminescence. Gist of method is development of curve of intensity of luminescence according to the exponential law in time on screen of cathode oscillograph. By varying the exponent of the converter, straight line produced on the screen, indicating identical nature of the law of attenuation of studied luminescence with the law of the converter. Submitted by Academician S. I. Vavilov, 9 Nov 1947.

PA 43/43T101

PA 48/49T98

TEOFILOV, P. P.

Dec 48

USSR/Physics
Luminescence
Polarization

"Polarized Luminescence," P. P. Tefilov, 40 pp

"Uspehi Fiz Nauk" Vol XXVI, No 4

Treats under: (1) Polarized luminescence as a result of anisotropy of the molecule, (2) observations on and measurements of polarized luminescence, (3) rotating depolarization and some problems of the theory of the liquid state, (4) concentrated depolarization and intramolecular exchange of energy of excitation, and (5) polarization of luminescence and structure of the molecule.

48/49T98

Dec 48

USSR/Physics (Contd)

molecule. Under (5), author discusses: (a) nature of elementary radiations and polarization diagrams, (b) polarized spectra and relative disposition of molecular oscillators, and (c) boundary polarization of luminescence and symmetry of the molecule.

48/49T98

FEOFILOV, P.

PA 48/49T93

USSR/Physics
Academy of Sciences
Luminescence

Dec 48

"Second All-Union Conference on Luminescence,"
P. Feofilov, 10 pp

"Uspekhi Fiz Nauk" Vol XXXVI, No 4

Session convened 17 - 22 May in Moscow. Discussed various problems of luminescence and illumination engineering. Vavilov summarized development of this science. Tarenin reported on "Quenching of Photoluminescence and Metastable States." Feofilova discussed polarized luminescence and structure of the molecule. Vavilov

48/49T93

also submitted a report on experimental observation of migrations of the exciting energy in fluorescent solutions.

48/49T93

ANDRONNIKOV, K.S.; BALAKOV, V.V.; BUZHINSKIY, A.N.; BURAGO, A.N.; VENTMAN, L.A.; VISHNEVSKIY, A.A.; VOLOSOV, D.S.; GASSOVSKIY, L.N., professor; GERSHUN, A.A., professor; YEL'YASHEVICH, M.A.; YEVSTROP'YEV, K.S.; GURVICH, M.M., professor; KOLYADIN, A.I.; KORYAKIN, B.M.; KURITSKIY, A.L.; PAPIYANTS, K.A.; PROKOF'YEV, V.K., professor; PUTSEYKO, Ye.K.; REZUNOV, M.A.; RITYN', N.E.; SAVOST'YANOVA, M.V., professor; SEVCHENKO, A.N.; SENNOV, N.I.; STOZHAROV, A.I.; FAYERMAN, G.P., professor; FEOFILOV, P.P.; TSAREVSKIY, Ye.N., professor; CHEKHMATAYEV, D.P.; YUDIN, Ye.F.; KAVRAYSKIY, V.V., professor; VAVILOV, S.I., akademik, redaktor

[Optics in military science] Optika v voennom dele; sbornik statei. Pod red. S.I.Vavilova i M.V.Savost'ianovoi. Izd. 3-e, zanovo perer. i dop. Moskva. Vol.2. 1948. 387 p. (MLRA 9:9)

1. Akademiya nauk SSSR. 2. Sostaviteli - sotrudniki Gosudarstvennogo Opticheskogo instituta (for all except Vavilov and Kavrayskiy)
3. Voenno-morskaya akademiya (for Kavrayskiy)
(Optics)

PA 36/49193

PEOFILOV, P. P.

USSR/Physics
Luminescence,
Molecular Structures

Jan/Feb 49

"Luminescence Polarization and Molecular Structure," P. P. Pefilov, State Opt Inst, 10 pp

"Iz Ak Nauk SSSR, Ser Fiz" Vol XIII, No 1

Approached study of molecular structure by investigating luminescence polarization, in particular, by: (1) polarization diagrams indicating spatial distribution of luminescence polarization, (2) polarization spectra, i.e., curves of observed polarization versus wave

36/49193

USSR/Physics (Contd)

Jan/Feb 49

length of exciting light, and (3) boundary values for degree of polarization, i.e., values observed when all known depolarizing factors were eliminated.

36/49193

117 AND 118 COVER										119 AND 120 COVER									
PROCESSES AND PROPERTIES INDEX										335,371									
<div style="position: absolute; top: 10px; left: 10px; font-size: 2em; font-weight: bold;">SA</div>										<div style="position: absolute; top: 10px; right: 10px; font-size: 2em; font-weight: bold;">A53</div> <div style="position: absolute; top: 10px; left: 10px; font-size: 2em; font-weight: bold;">J</div>									
<p>166. New method of investigating the processes of growth and decay of luminescence. N. A. TOLSTOI AND P. P. FROHLIN. <i>Izv. Akad. Nauk, SSSR, Ser. Fiz.</i>, 43 (No. 2) 211-17 (1949) <i>In Russian</i>.</p> <p>A general method of studying relaxation phenomena has been worked out using a specially designed sweep circuit equipped with a time-base following approximately the law of the process studied. When the method is applied to the processes of growth and decay of luminescence, the value of the parameter τ (from the exponential time-function $x = x_0 e^{-t/\tau}$) can be measured with an accuracy of about 1% (under favourable conditions).</p> <p style="text-align: right;">F. LACHMAN</p>																			
ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION										6-EXTENDED TABLE									
FROM SYNOPTIC										FROM GENERAL									
SYNOPTIC										GENERAL									
SYNOPTIC										GENERAL									

FEOFILOV, P. P.

USSR/Physics
Relaxation Phenomenon
Oscillographs

May 49

"New Method of Studying Processes," N. A. Tolstoy, P. P. Feofilov, State Optical Inst, 10 pp

"Zhur Eksper i Teoret Fiz" Vol XIX, No 5

Describes new oscillographic method of studying relaxation of physical processes, based on the use of nonlinear graphing. Method is applicable to processes occurring in the time interval 10^{-6} - 10^{-1} sec. In the case of relaxation taking place according to the exponential law ($\exp - t/\tau$), value of τ may be determined after several seconds. Generalization of method, called "the method of partial time," is applicable to processes in which relaxation occurs according to any law. Submitted 6 Nov 48.

PA 46/49T99

FEOFILOV, P. P.

USSR/Physics - Dielectric Constants
Polarization, Relaxation

Dec 49

"Application of the New Method for Studying Relaxation Processes to a Study of Relaxation Polarization in Dielectrics," G. I. Skanavi, N. A. Tolstoy, P. P. Feofilov, K. I. Lebedev, Phys Inst imeni Lebedev, Acad Sci USSR, 9 pp

"Zhur Eksper i Teoret Fiz" Vol XIX, No 12

To study relaxation polarization in dielectrics of titanium dioxide with small additions of oxides of metals belonging to the second group of the periodic table (this group gives very high values of ϵ in the region of low frequencies), one employs the oscillographic method of studying, by electrical square-wave impulses through ohmic resistances, the charge and discharge of the condensers containing the dielectric under study. Here a simple exponential development of the process in time is employed, as well as more complicated ones. Shows charge and discharge processes of the condenser with the dielectric under study have a complex character differing from the exponential. Equivalent circuit schemes are found for the dielectrics under study. Parameters of these schemes are determined experimentally. Calculation of these equivalent schemes permits one to obtain the function of current drop with time in each studied dielectric with calculated constants and thus to evaluate values of initial currents. Submitted 23 Jun 49

PA 152T87

FEOFILOV, P. P.

USSR/Physics
Phosphors
Luminescence

Jan 49

"A Study of the Initial Stages of Luminescence and Extinguishing in Zinc-Sulfide Phosphors With the Aid of an Oscillographic Phosphoroscope," V. A. Arkhangel'skaya, A. M. Bonch-Bruyevich, N. A. Tolstoy, P. P. Feofilov, 4 pp

"Dok Ak Nauk SSSR" Vol LXIV, No 2

Partial results obtained during study of crystalline phosphors by the "partial time" method. Method allows studies to be conducted in the time interval 10^{-5} - 10^{-1} seconds, suitable for investigating the little-studied initial stages of extinguishing, and the completely unstudied stages of crystalline phosphors bursting into luminescence. Submitted 5 Nov 48.

PA 25/49T99

FEOFILOV, P. P.

USSR/Physics
Optics
Colloids

Jun 49

"Certain Electro-Optical Phenomena in Colloids," N. A. Tolstoy, P. P. Feofilov,
State Opt Inst, 4 pp

"Dok Ak Nauk SSSR" Vol LXVI, No 4

Studied electrical double refraction and electrical dichromism of colloids using (1) optical arrangement with a Kerr cell, (2) voltage sources, changing according to a special law, (3) obdurator, and (4) photoelement, amplifier, and cathode oscilloscope. Tested water solutions of bentonite, benzopurpurin, graphite, anisaldazin, n-azoxyanisole, and other liquid crystals. Submitted by S. I. Vavilov, 25 Mar 49.

PA 46/49T89

FEOFILOV, P. P.

PA 52/49T105

USSR/Physics

Photoconductivity

Bismuth Compounds

May 49

"Initial Stages of Relaxation in the Photoconductivity of Bi_2S_3 ," D.B. Gurevich,
N. A. Tolstoy, P. P. Feofilov, 4 pp

"Dok Ak Nauk SSSR" Vol LXVI, No 3. Submitted by Acad S. I. Vavilov, 25 Mar 49.

CA

3

Some particularities of the flare-up of ZnS:Mn phosphor. N. A. Tolstol and P. P. Kuznetsov. *Doklady Akad. Nauk S.S.S.R.* 248, 248 (1980). In excitation by a Hg-vapor lamp, the blue (Zn) band of the ZnS:Mn phosphor shows, in oscillographic phosphoroscopes, normal growth of intensity, but the yellow (Mn) band first flares up to a peak, from which it rapidly falls to the stationary intensity. In samples heated 15 min. at 900°, under a NaCl flux, with $1 \times 10^{-3} - 1 \times 10^{-4}$ g. Mn/g., the peak was most pronounced at 1×10^{-3} g./g., and was still observable with 5×10^{-4} and 5×10^{-5} g./g. The effect is thus most marked at activator concns. at which the competition between the blue and the yellow emission is most pronounced. The height of the peak (relative to the stationary intensity) increases with the intensity of excitation, and with the length t_0 of the (equal) periods of illumination and darkness (between 0.01 and 0.4 sec.). On cooling to liquid-air temp., the effect disappears in samples in which it was present at room temp., whereas samples with too high or too low Mn contents show no new particularities. On heating to above room temp., the relative height of the peak first increases, then falls, along with a general increase of the stationary brightness, with a distinct min. appearing between the peak and the level branch. Further increase of the temp. brings the curve back to normal, without peak or min. The peak phenomenon is thus confined to a range close to room temp. By exponential spreading, the max. of the peak is found to be attained within a few millisees. from the beginning

of the excitation; this time is practically independent of t_0 , and decreases with increasing intensity of excitation. The peak can be made visually observable with the aid of a rotating disk coated with the phosphor on its periphery; the brightness of the excited luminescent spot, if viewed through a red filter, first increases, then decreases with increasing speed of rotation, whereas the contrary appears through a blue filter. The decay of the yellow emission also presents peculiarities. It is very fast in the beginning of the dark period, then is slowed down almost to a standstill, then proceeds at the normal rate. This effect is most pronounced with 5×10^{-4} g. Mn/g., and is absent altogether above 5×10^{-3} or below 1×10^{-4} g./g. Again the blue emission decays normally throughout. The phenomenon of the flare-up can be interpreted in agreement with the representations of Levshin (C.A. 42, 3386c), by assuming that at the end of a dark period of sufficient length t_0 , the levels of Mn have become empty, as a consequence of which electron transitions to these levels are uninhibited in the 1st stages of excitation, but become stationary as soon as the levels have become "plugged" with electrons. This mechanism accounts satisfactorily for all the effects observed, except for the disappearance of the flare-up at too low Mn concns., and for the min. observed at moderately high temps.

N. Thon

FEOFILOV P. F.

Relaxation of the photoconductivity in semiconductors. D. B. Gurevich, N. A. Tolstoi, and P. F. Feofilov. Zhur. Ekspil. Teoret. Fiz. 20, 769-82 (1950).-- The kinetics of the photocond. $\Delta\sigma$ were investigated by oscillography, in illumination with rectangular light impulses (mostly of a period ~ 0.04 sec.), with rectangular voltage impulses of a frequency very considerably greater than that of the light impulses, and with exponential sweep of the oscillograph screen. The exptl. relaxation curves are represented in the coordinates $\theta = \Delta\sigma / (d\Delta\sigma/dt) = dt/d(\ln\Delta\sigma)$ vs. time t ; the magnitude θ signifies an "instantaneous relaxation time," i.e. the relaxation time τ of the exponential $e^{-t/\tau}$ which approximates the exptl. curve at any point corresponding to the moment t . In other words, θ expresses at any given moment the "sp. slowness" of the relaxation, i.e. the reciprocal of the rate of fall of σ relative to the distance from equil. For simple exponential processes, θ coincides with the relaxation time τ , i.e. exponential processes are characterized by a const. θ . The rate law of the relaxation can be deduced from the shape of the $\theta(t)$ plots. (1) For Bi_2S_3 (thin layer on glass), θ is a linear function of t at any temp. between 189 and 326°K., i.e., $\theta = A + Bt$, or, with $A = 1/\alpha$, $B = 1/\alpha$, and after integration, $\Delta\sigma = \Delta\sigma_0 / (1 + \alpha t)^{\alpha}$, where $\Delta\sigma_0$ = initial photocond. This hyperbolic law of the decrease of $\Delta\sigma$ is analogous to the Becquerel law in phosphorescence. Between 189 and 326°K., α varies from 0.039 to 0.23, following $\alpha = \alpha_0 e^{uT/4}$, with $u = 6.2 \times 10^{-3}$ e.v., and α_0 (dimensionless) = 1.23×10^{-3} . The parameter α is roughly independent of the temp. and is = 1200 sec.⁻¹ at max. illumination E . The plot of the stationary $\ln\Delta\sigma$ vs. $\ln E$ is linear. At const. temp., α is approx. proportional to $E^{1/\beta}$, with $\beta = 3-4$, and the variation of $\Delta\sigma$ with E obeys approx. the same law. This gives the simple relation $\alpha = \alpha E^{-\beta} + \rho \Delta\sigma$. The growth of $\Delta\sigma$ during illumination follows a complex nonexponential law; its

temps. the growth of $\Delta\sigma$ is exponential. The exponent α in the decay law increases with temp., particularly rapidly between ~ 150 and $\sim 200^\circ$; $a - a_0 = k_3 E^{1/2}$, with k_4 falling with rising temp. In the growth law $\Delta\sigma = \Delta\sigma_0 (1 - e^{-t/\tau})$, $1/\tau = k_4 E^{1/2}$, with k_4 falling with rising temp. With further rising temp., the growth curve deviates increasingly from the exponential. The values of a and $1/\tau$ are of the same order of magnitude. Above 200°C ., both the growth and the decay are exponential, with τ independent of E but varying with the temp. according to $\tau \sim e^{-u/KT}$ with $u = 0.45$ e.v. (7) Photoconductors can be divided into 2 categories; (I) "exponential" photoconductors, characterized by a stationary $\Delta\sigma_0 \sim E$, a dark cond. roughly 10-100 times the photocond., exponential growth and exponential decay of $\Delta\sigma$ with the same τ in both instances, and independent of E ; the temp. variation of $1/\tau$ is analogous to the temp. variation of the dark cond., and of the form $e^{-u/KT}$ in both instances; examples of this category are Cu_2O , and CdS at higher temps. (II) "Hyperbolic" photoconductors, characterized by $\Delta\sigma_0 \sim E^{1/2}$, a relatively low dark cond., a decay following $\Delta\sigma = \Delta\sigma_0 / (1 + at)^\alpha$, with α independent of E , and $a \sim E^{1/2}$; the growth can be exponential, in which case $1/\tau \sim E^{1/2}$. Examples are Bi_2S_3 , Tl_2S , Se , InSe , and CdS at low temps. Possible deviations of certain properties (except the hyperbolic decay) from this simple scheme are viewed as complications only.

N. Thon

FEOFILOV, P. P.

USSR/Physics - Phosphore Photoconductivity

Nov 50

"Comparison of Photoconductivity Relaxation," D. B. Guravich, H. A. Tolstoy,
P. P. Feofilov

"Zhur Eksper i Teoret Fiz" Vol II, No 11, pp 1039-1046

Compares experimental laws governing kinetics of photoconductivity with experimental laws governing kinetics of luminescence. Establishes parallelism of these laws.

Calculates luminescence relaxation of cadmium sulfide from its photoconductivity relaxation. Submitted 4 Apr 50.

PA 169T107

C 4

New method of investigation of relaxation processes and its application to the study of some physical processes. N. A. Tolstol and P. P. Frolov. *Uspekhi Fiz. Nauk* 41, 44-107(1950).--Review, particularly of the authors' own work on rectilinear oscillography of the decay of phosphorescence, photocond., etc. 33 references. N. Thon

FEOFILOV, P. P.

1 Mar 50

USSR/Physics - Luminescence
Conductivity, Photo-

"Luminescence and Photoconductivity of Cadmium Sulfide," D. V. Gurevich, N. A. Tolstoy, P. P. Feofilov

"Dok Ak Nauk SSSR" Vol LXXI, No 1, pp 29-32

Investigated kinetics of luminescence and photoconductivity, carried out by method of "taumeter" (tau— life span or duration) on CdS monocrystals. Mathematically discusses phosphorescence relaxation and so-called "nonlinear" photoconductivity. Taumeter was described in ZhETF 19, 421, 1949, and "Iz Ak Nauk SSSR, Ser Fiz," 13, 211, 1949, by Tolstoy et al. Submitted 3 Jan 50 by Acad S. I. Vavilov.

PA 165T75

FEOFILOV, P. P.

A. N. Terenin, acad., and P. P. Feofilov. The greatest Soviet scientist-optician.
(refers to S. I. Vavilov) P. 111

SO: Vestnik, Messenger of the Acad. of Sci., USSR. 21, 3, (1951).

WE.

*Materials & Subsidary
Techniques*

333.37 2106
The Third All-Union Conference on Luminescence and
the Applications of Luminescent Materials. - P. Fezhloy
(Uspekhi fiz. Nauk, Nov. 1951, Vol. 45, No. 5, pp.
445-457) Report on a conference held in Moscow, June
15-21, 1951. Summaries of 45 papers are given

FEOFILOV, P. P.

USSR/Physics - Semiconductors

Jan/Feb 52

"Kinetics of Photoconductivity and Kinetics of Phosphorescence," N. A. Tolstoy, P. P. Feofilov

"Iz Ak Nauk SSSR, Ser Fiz" Vol XVI, No 1, pp 59-69

Separates relaxation of photocond into 2 basic types: hyperbolic and exponential. Considers non-linear photoresistance with relatively low dark-current cond, characterized by bimol recombination mechanism, as belonging to the hyperbolic type; while linear photoresistance with high dark-current cond belongs to the exponential type.

218187

1. FEOFILOV, P. P.
2. SSSR (600)
4. Vavilov, Sergei Ivanovich, 1891-1951
7. New developments in the theory of light ("Microstructure of light.")
S. I. Vavilov. Reviewed by P. P. Feofilov.
Usp. fiz. nauk 46 No. 1, 1952
9. Monthly List of Russian Accessions, Library of Congress, February 1953. Unclassified.

FEOFILOV, P. P.

PA 236T76

USSR/Physics - Optics, Book Review

Sep 52

"New Paths in the Development of Theory of Light
(Review of S. I. Vavilov's book, 'Microstructure of
Light')," P. P. Feofilov

"Uspekhi Fiz Nauk" Vol 48, No 1, pp 3-24

Praises Vavilov's work, describing visual observations of quantum fluctuations of light, principles of superposition and "nonlinearity" in optics, micro-optical analysis of interference, Vavilov-Cherenkov effect, microoptics of absorbing media, inductive resonance, and migration of energy. Expects book to be a perfect manual for future generations.

236T76

FEOFILOV, P. P.

235T93

USSR/Physics - Luminescence Extinction 21 Jul 52

"Problem of Extinction of Luminescence of Rubin
($\text{Al}_2\text{O}_3\cdot\text{Cr}$)," N. A. Tolstoy, P. P. Feofilov

"Dok Ak Nauk SSSR" Vol 85, No 3, pp 551-554

Presents certain data on concn, temp, and spectral dependences of curves of extinction (and lighting) of luminescence of synthetic ruby, from which it follows that the mechanism governing luminescence in rubies is rather complex. Submitted by Acad A. N. Terenin 26 May 52.

235T93

FEOFILOV, P. P.

PA 240T97

USSR/Physics - Luminiscence

21 Dec 52

"Absorption and Luminiscence Spectra of Triphenylmethane Dyes," P. P. Feofilov and I. G. Fayerman

"IAN SSSR" Vol 87, No 6, pp 931-934

Compare absorption and emission spectra of triphenylmethane dyes in order to verify that luminescence spectra should be narrower than absorption band. Results are tabulated. Besides fluorescence, the dyes emit for a few seconds a greenish afterglow in a wave-length shorter than the exciting light. Presented by A. N. Terenin. Received 21 Oct 52.

240T97

PA 254T61

FEOFILOV, P. P.

USSR/Electronics - Radio Location, History Jan 53

"Unknown Work by S. I. Vavilov on Locating Hidden
Radios," P. P. Feofilov and I. A. Shlyakhter

Usp Fiz Nauk, Vol 49, No 1, pp 147-154

Describes an unpublished work by S. I. Vavilov con-
cerning the location of enemy radio transmitters
during World War I.

254T61

FEOTILOV, P.P.

Polarization of the luminescence of F-centers in crystals of alkali halides. P. P. Feofilov. *Doklady Akad. Nauk S.S.S.R.* 92, 743-745 (1954) (Engl. translation issued as *U.S. Atomic Energy Comm. NSF-tr-208*, 4 pp. (1954)); cf. *C.A.* 48, 7439b. The study was made with large LiF and NaF single crystals. The crystals were colored by x-irradiation. The absorption spectrum of NaF presented peaks at about 340 and 510 mμ. The absorption spectrum of LiF was almost identical with that reported by Klick (*C.A.* 45, 10059c). The red-orange luminescence of NaF was excited with a spectrum practically unchanged over the entire region of spectral absorption of the crystal and was quite intense with excitation in both absorption bands. Excitation of x-rayed LiF by 430-mμ light yielded a bright orange luminescence, while excitation by 385-mμ light resulted in bright green luminescence. Duration of luminescence was much less than 10 microsec. The luminescence of x-rayed LiF and NaF single crystals excited by linearly polarized light was partially polarized. For NaF, the polarization changed sign with transition of excitation from the vicinity of the long-wave absorption band to the short-wave band. Because the emission spectrum was almost unchanged, it was concluded that in NaF both bands must be ascribed to centers of the same type. It was assumed that the long-wave oscillators were distributed along the 6 C_2 axes which pass through the centers of the opposite edges of the elementary cube. The circular oscillators describing the short-wave absorption were arranged in the (110) planes of the rhombododecahedron. Calcns. were made which resulted in excellent agreement for the azimuthal dependence of the degree of polarization. The abs. values calcd. differ somewhat from exptl. values because the anisotropy of the oscillators was not complete. Spontaneous polarization under excitation by natural light was observed in plates cut parallel to the (110) plane. Calcns. were in full agreement with the exptl. data. H. L. Jr.

PEOFILOV, P. P.

USSR/Physics - F-Centers Crystals

21 Sep 53

"Photochemical Conversion of F-Centers in KCl Crystals at High Temperatures," A.A. Chatalov, Kiev State Univ in Shevchenko.

DAN SSSR, Vol 92, no 3, pp 549-552

Studies effect of light at high temps on absorption of KCl crystals containing F-centers. Used method of S.A. Artsybashev (Trudy Fiz Inst Lebedev, Vol 2, No 3 (1938)). Plots the variation of X-band, obtained at const irradiation of crystal and decaying after discontinuance of irradiation. Considers X-centers as a primary formation of colloidal coagulation of several or, most probable, 2 F-centers. Presented by Academician A.N. Terenin 21 Jul 53.

268T95

FEOFILOV, P. P.

USSR/Physics - Luminiscence,
F-Centers

21 Sep 53

"Polarized Luminiscence of F-Centers in CaF_2 ," P.P.
Feofilov

DAN SSSR, Vol 92, No 3, pp 545-548

Revealed intensive red luminiscence of CaF_2 crystals, dyed by additive or subtractive method. This luminiscent excitation spectrum coincides with the absorption spectrum of F-centers and therefore may be ascribed to F-centers. The latter may be interpreted as electrons localized in holes of halide ions. Presented by Acad A.N. Terenin 20 Jul 53.

268T94

USSR/Minerals - Spectral analysis

Card 1/1 Pub. 43 - 91/97

Authors : Feofilov, P. P., and Kuznetsova, L. A.

Title : Spectral luminescence method of determining the Cr content in synthetic rubies

Periodical : Izv. AN SSSR. Ser. fiz. 18/2, page 297, Mar-Apr 1954

Abstract : A method was developed for quantitative spectral determination of chromium in natural and synthetic rubies. A study of spectral, kinetic and polarization characteristics of the luminescence of rubies with various Cr contents showed that a change in Cr concentration affect mostly the luminescence spectra. The Cr ions which are isomorphically included in the crystalline lattice of corundum (Al_2O_3), determine the red color intensity and the luminescence of natural and synthetic ruby crystals.

Institution :

Submitted :

FEOFILOV, P. P.

USSR/ Physics - Spectral analysis

Card 1/1 Pub. 43 - 27/62

Authors : Feofilov, P. P.

Title : Spatial distribution of radiation and the nature of coloration centers in ion crystals

Periodical : Izv. AN SSSR. Ser. fiz. 18/6, 688-689, Nov-Dec 1954

Abstract : It was established that the luminescence of coloration centers forming during photochemical or additive coloring of CaF_2 , NaF and LiF monocrystals and excited by a linear-polarized light is partially polarized. The observed absolute values of the polarization indicate the high anisotropy of the radiation centers of the coloring matter. The relation between the degree of polarization and the relative orientation of the crystal and the exciting electrical vector, is discussed. Graph.

Institution:

Submitted :

FD 425

USSR/Physics - Color centers

Card 1/1

Pub. 147-11/16

Author : Feofilov, P. P.

Title : Anisotropy of the radiation of color centers in crystals of cubic syngony

Periodical : Zhur. eksp. i teor. fiz. 26, 609-623, May 1954

Abstract : On the basis of a classical oscillator model the author calculates the azimuthal dependence of the degree of polarization of luminescence of the color centers in cubic crystals. Conducts the computation for the cases where the elementary oscillators are oriented along the basis of symmetry of the fourth, third, and second orders and for the excitation of luminescence in various regions of the absorption spectrum. Comparison of the computed and experimental data permits one to establish the character of the orientation of the color centers in various cases investigated and to express assumptions concerning the nature of the centers.

Institution :

Submitted : September 4, 1953

"APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5

APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5"

USSR/Physics - Crystallography

Card 1/1 Pub. 22 - 17/56

Authors : Feofilov, P.P.

Title : About orientation of the Eu^{++} ions in the lattice of crystal CaF_2

Periodical : Dok. AN SSSR 99/5, 731-733, Dec. 11, 1954

Abstract : Results of the fluorescence-polarizing method, applied to CaF_2 crystals, for determining the orientation of Eu^{++} ions on the crystal lattices are described. Four references: 3-USSR (1934-1954) Graph; Photodiagrams.

Institution:

Presented by: Academician A.N. Terenin, August 6, 1954

FEOFILOV P.P.

K-5

Category : USSR/Optics - Physical Optics

Abs Jour : Ref Zhur - Fizika, No 2, 1957, No 4963

Author : Feofilov, P.P.

Title : Experimental Determination of the Nature of the Elementary Radiators of the Eu^{3+} Ion in CaF_2 .

Orig Pub : Dokl. AN SSSR, 1954, 99, No 5, 975-958

Abstract : The method of polarization luminescence diagrams, proposed by S.I. Vavilov (Zh. eksperim i teor, Fiziki, 1940, 10, 1363), for the determination of the nature (multi-polarity) of elementary radiators, is applicable to the investigation of the luminescence of Eu^{3+} in single crystals of CaF_2 -Eu. Employing data obtained in an earlier work (Referat Zh. Fizika, 1956, 5304) with respect to the character of the orientation of the elementary radiators of the Eu^{3+} ion (orientations along the third-order symmetry axes) the author computed the dependence of the intensity (J) and polarization (P) of the luminescence -- excitation through the face parallel to (100), observation through the face parallel to (110) -- on the angle of rotation (γ) of the electric vector exciting

Card : 1/2

Category : USSR/Optics - Physical Optics

K-5

Abs Jour : Ref Zhur - Fizika, No 2, 1957, No 4963

the linearly-polarized light. A calculation carried out for four types of elementary dipole radiators (electric and magnetic linear π_e and π_m and circular σ_e and σ_m) -- shows that the investigation of the functions $J(\eta)$ and $P(\eta)$ leads to an unambiguous conclusion concerning the multipolarity of the radiators (under the assumption that the absorption of light is carried out by linear electric oscillators). In many cases there is no need for a full investigation of the functions $J(\eta)$ and $P(\eta)$, and the determination of the nature of the radiators can be reduced to a qualitative observation of the sign of P and of the character of the function $J(\eta)$. The use of this method has made it possible to ascertain, that among the most intense lines in the luminescence spectrum of the Eu^{3+} ion in CaF_2 there are lines corresponding to all four possible types of dipole radiators (π_e , π_m , σ_e , and σ_m).

Card : 2/2/

FEOFILOV, P. P., Doctor of Physico-Mathematical sciences

"The Polarized Luminescence of Cubic Crystals" paper presented at the Conference on Molecular Luminescence and Luminescent Analysis, MINSK from 20 to 25 June 1955.

PEOFILOV, Petr Petrovich,

Academic degree of Doctor of Physio-Mathematical Sciences, based on his defense, 29 June 1955, in the Council of State Order of Lenin Optical Inst imeni Vavilov, of his dissertation entitled: "Polarized luminescence of atoms, molecules, and crystals."

Academic degree and/or title: Doctors of Sciences

SO: Decision of VAK, List no. 4, 25 February 1956, Byulleten' MVO SSSR, No. 1, January 1957, Moscow, pp. 14-24, Uncl.
JPRS/NY-440

FD-2982

USSR/Physics - Absorption band

Card 1/1 Pub. 146 - 23/28

Author : Grechushnikov, B. N.; Feofilov, P. P.

Title : Oscillatory structure in the absorption spectrum of rubidium at
minus 190° C

Periodical : Zhur. eksp. i teor. fiz., 29, September 1955, 384

Abstract : S. I. Pekar (ibid., 22, 641, 1952) and M. A. Krivoglaz (co-author
S. I. Pekar, Trudy Instituta fiziki AN Ukr. SSR, 4, 37, 1954)
showed theoretically that the form of the absorption band for ad-
mixtures in dielectrics can be represented under definite assump-
tions in the form of the product of two frequency functions, the
first factor representing the smooth bell-shaped curve and de-
scribing the general contour of the absorption spectrum and the
second factor representing a rapidly oscillating function and de-
scribing the so called structure of the absorption band. B. N.
Grechushnikov (DAN SSSR, 99, 707, 1955) investigated the absorp-
tion spectrum of rubidium at the temperature of liquid nitrogen
(the earlier study of the influence of temperature was by K. Gib-
son, Phys. Rev., 8, 38, 1916). The present writer obtains greater
detail. Five references: e.g. Krishnan, Proc. Ind. Soc., 26, A,
6, 450, 1947.

Institution : Institute of Crystallography, Academy of Sciences USSR

Submitted : May 19, 1955

FEOFILOV, P. P.

USSR/Physics - Conferences

Card 1/1 Pub. 118 - 5/8

Authors : Neporent, B. S. and Feofilov, P. P.

Title : The 9th conference on spectroscopy

Periodical : Usp. fiz. nauk 55/3, 443-452, Mar 1955.

Abstract : A description of the work conducted at the 9th conference on spectroscopy is presented. The conference took place at Tartu (Estonia) during July 5 to 11, 1954. The conference was divided into 3 separate sections: 1. molecular spectroscopy; 2. general and atomic spectroscopy; and 3. spectral analysis. Reports on molecular spectroscopy, read by many scientists, are outlined.

Institution:

Submitted :

FEOFILOV, P. P.

K-5

Category : USSR/Optics - Physical Optics

Abs Jour : Ref Zhur - Fizika, No 2, 1957, No 4959

Author : Feofilov, P. P.

Title : Concerning the Theory of Polarization of Resonant Radiation and Fluorescence of Atoms and Diatomic Molecules.

Orig Pub : Dokl. AN SSSR, 1955, 104, No 6, 846-849

Abstract : The degree of polarization $P(J)$ and the degree of circularity $C(J)$ of fluorescence is calculated for plane-polarized or circularly-polarized light respectively. The population of the upper level is determined by the probability of the dipole transition $A_{J_2 M_2}^{J_1 M_1}$ in the former case and $A_{J_2 M_2}^{J_1 M_1}$ in the latter. The intensity of the π and σ components is determined by the same population and by the probability of the reverse transition $M \rightarrow M$ (π) and $M \rightarrow M \pm 1$ (σ). This yields $P(J) = \frac{I_\pi - I_\sigma}{I_\pi + I_\sigma}$ and analogously $C(J) = \frac{I_+ - I_-}{I_+ + I_-}$ where I_k is the intensity of the components with right and left-hand circular polarization. The results are given for the transitions $J_1 \rightarrow J_2 \rightarrow J_3$ (for all nine combinations, satisfying the selection rules) as functions of J_1 . The equations obtained

Card : 1/2

VAVILOV, S.I.; LEBEDEV, A.A., akademik; TOPCHIEV, A.V., akademik; TRENIN, A.N., akademik; LANDSBERG, G.S., akademik; VUL, B.M.; KRAVETS, T.P. [deceased]; LEVSHIN, V.L.; FEOKILOV, P.P.; GALANIN, M.D.; KUZNETSOV, I.V.; VAVILOV, V.S.; GUROV, K.P., redaktor izdatel'stva; KISELEVA, ALAL, tekhnicheskiy redaktor

[Collected works] Sobranie sochinenii. Moskva, Izd-vo Akademii nauk SSSR. Vol.4. [Experimental foundation of the theory of relativity. On "warm" and "cold" light. The eye and the sun. Popular scientific articles and reviews] Eksperimental'nye osnovaniia teorii otnositel'nosti, O "teplom" i "kholodnom" svete, Glaz i solntse, Nauchno-populiarnye i obsornyie stat'i. 1956. 469 p. (MLRA 9:8)

1. Chlen-korrespondent AN SSSR (for Vul, Kravets)
(Physics)

"APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5

APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5"

"APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5

APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5"

EE OF 11 019 P.P.

✓ Vibration structure in the absorption spectrum of ruby.
B. N. Grechushnikov and P. P. Feofilov. *Soviet Phys.*
JETP 2, 330 (1956) (Engl. translation).— See C.A. 50
2205a. *Phys.*

3

TEOFILOV, P.P.

TEOFILOV, P.P.

Polarized luminescence of cubic crystals. Izv.AN SSSR Ser.fiz.no.5:
552 '56. (luminescence) (MIRA 9:9)

FEOFILOV, P.

Sergei Ivanovich Vavilov, March 24, 1891--Jan 25, 1951. Opt.1
spektr. 1 no.2:107-112 Je '56. (MLRA 9:11)
(Vavilov, Sergei Ivanovich, 1891-1951)

FEOFILOV, P. P.

B-5

• USSR/Crystals.

Abs Jour : Referat Zhur - Khimiya, No 6, 1957, 18340

Author : P.P. Feofilov.

Title : Nature of Elementary Oscillators and Polarization of
Photoluminescence in Cubic System Crystals.

Orig Pub : Optika i spektroskopiya, 1956, 1, No 2, 131-142

Abstract : The possibility of an experimental determination of the multipole order of elementary oscillators of luminescence centers orientated along different axes of symmetry of cubic system crystals is discussed in general. The theoretical computation of the dependence of the luminescence intensity and polarization on the reciprocal position of the observation direction and the electrical vector of the exciting light is carried out for three possible cases of the center orientation (along the axes of symmetry of the 2nd, 3rd and 4th order) and for all the possible combinations of dipole oscillators

Card 1/2

- 97 -

State Optical Inst.

USSR/Crystals.

B-5

Abs Jour : Referat Zhur - Khimiya, No 6, 1957, 18340

(electrical and magnetic, linear and circular). For the same oscillator combinations, the corresponding dependences are computed for randomly distributed centers (isotropic solutions). The computed dependences show that in the majority of cases, the multipole order of absorbing and radiating oscillators can be determined uniquely. The method is applicable in presence of polarized luminescence in investigated centers. Experimental data confirming the usefulness of the method are quoted.

Card 2/2

- 98 -

FEVFILOV, P.
ARKHANGEL'SKAYA, V.; TOLSTOY, N.; ~~FEVFILOV, P.~~

Fifth Conference on Luminescence (crystal phosphors). Opt.
1 spektr. 1 no.6:813-820 0 '56. (MLRA 9:12)

(Tartu--Luminescence--Congresses)

FEofilov, P. P.

Category: USSR / Physical Chemistry - Crystals

B-5

Abs Jour: Referat Zhur-Khimiya, No 9, 1957, 29743

Author : Feofilov P. P.

Inst : not given

Title : Anisotropy of Luminescent Coloration Centers in CsI-Tl Crystals

Orig Pub: Optika i spektroskopiya, 1956, 1, No 7, 952-954

Abstract: Investigation of anisotropy of radiation of coloration centers which arises on irradiations of crystals of CsI-Tl with ultraviolet, x-ray or gamma radiation. As a result of study of azimuth dependence of luminescence polarization it was found that luminescence polarization depends strongly upon the relative orientation of exciting electric vector and the crystallographic axes, and that the anisotropic luminescent coloration centers are oriented along 4-th order symmetry axes of CsI crystals. Study of polarization spectrum and of dependence of the degree of polarization upon wave length of radiation, shows that degree of polarization is greatly dependent on radiation wave length. This is attributed to the fact that the radiation band

Card : 1/2

-33-

Category: USSR / Physical Chemistry - Crystals

Abs Jour: Referat Zhur-Khimiya, No 9, 1957, 29743

B-5

constitutes, apparently, a superposition of at least two bands, corresponding to different types of coloration centers. A center model is proposed: the center consists of Tl^{+} ion, isomorphously replacing a Cs^{+} ion and associated with another defect (Tl^{+} ion or a hole), which is also localized within the cation point.

Card : 2/2

-34-

Feofilov, P. P.

USSR/Optics - Physical Optics

K-5

Abs Jour : Referat Zhur - Fizika, No 5, 1957, 12923

Author : Tishenko, G.A., Feofilov, P.P.

Inst : -

Title : Luminescence of Color Centers in Crystals of Fluorite

Orig Pub : Izv. AN SSSR, ser. fiz., 1956, 20, No 4, 482-487

Abstract : An investigation was made of the contours of the absorption band and of the luminescence of the color centers in artificial crystals of fluorite. The bell-shaped form of the bands is satisfactorily described by the theory of impurity absorption (Pekar). The Stokes shift of the bands, calculated with formulas of this theory using data on the half widths of the bands, is also in agreement with the experimental data. The method of polarization of the luminescence diagrams was used to establish the electric dipole character of the elementary oscillators, describing the absorption and radiation. It is proposed that there

Card 1/2

USSR/Optics - Physical Optics.

K-5


Abs Jour : Ref Zhur - Fizika, No 5, 1957, 12923

exists an infrared luminescence of complex (double) color
centers in crystals of alkali-halide salts.

Card 2/2

"APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5



APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5"

USSR/ Physics - Scientific conferences

Card 1/1 Pub. 118 - 6/7

Authors : Naporent, B. S., and Feofilov, P. P.

Title : The Fourth Luminescence Conference (Molecular Luminescence and Luminescence Analysis)

Periodical : Usp. Fiz. nauk, 58/1, 151-164, Jan 1956

Abstract : The fourth conference on luminescence is recounted. The conference, which took place at Minsk, BSSR, was called by the Acad. of Scs. of the BSSR, Physical Institute imeni P. N. Lebedev, the Acad. of Scs. of the USSR and the Optical Institute imeni S. I. Vavilov. More than two hundred (200) scientists of various fields participated in the conference. Possibilities for studying intermolecular processes through the observation and analysis of luminescence were discussed.

Institution :

Submitted :

FEOTILOV, P.P.

Nature of the color of green fluorite
Zapata, Mexico. Observed as a
thin film on a glass plate. The
color is green and translucent.
The film is called as "fluorite" and is
observed at -100°. The partial color
spectrum will change from
green to blue. The color is
observed in the blue region
of the spectrum. The color is
observed in the blue region
of the spectrum.

"APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5

(1870)

W. B. (a)

APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5"

FEOFILOV, P. P.

USSR/Physical Chemistry. Crystals.

B-5

Abs Jour: Ref Zhur-Khimiya, No 5, 1957, 14530

Author : I. V. Stepanov, P. P. Feofilov

Inst : -

Title : On Two Types of Luminescence Spectra of Rare Earths in Artificial Fluorite Crystals

Orig Pub: Dokl. AN SSSR, 1956, 108, No 4, 615-618

Abstract: Monocrystals of synthetic CaF_2 , activated by fluorides of rare earths (TR: Pr, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tu; $\sim 10^{-4}\text{g/g}$) yield two types of luminescence spectra (L), which differ sharply in number and position of the lines. Type II (anomal) is observed in crystals, grown in reducible conditions (addition to the batch of 10^{-3}g/g of graphite powder). Type I (normal) is prepared after heating the ready crystals in an open hearth at a temperature of $900-1000^\circ$. The (L) spectra are independent of excitation and do not change during damping. The absorption spectra of types I and II differ from one another.

Card 1/2

USSR/Physical Chemistry. Crystals.

B-5

Abs Jour: Ref Zhur-Khimiya, No 5, 1957, 14530

Abstract: The centers of both types can coexist in a crystal. The conclusion was made that in both cases TR are present in the form of ions of type Tr^{+3} , but the spectra depend on the closest environment. Crystals of type I are formed as a result of the substitution of one of the eight F^- ions surrounding TR^{+3} by an O^{2-} ion. The nature of substitution for type II was not established. In natural CaF_2 crystals, heated in an open hearth at $600-700^\circ$, the brilliance of (L) increases and spectral lines appear of type II (Er, Dy) and I (Sm). Heating at $900-1000^\circ$ leads to a second change of the spectrum: the spectra of all TR becomes identical with the type I spectrum of synthetic CaF_2 -TR.

Card 2/2

FEOFILOV, P.P.

USSR / Physical Chemistry. Crystals.

B-5

Abs Jour : Ref Zhur - Khimiya, No 8, 1957, 25922.

Author : V.A. Arkhangel'skaya, P.P. Feofilov.

Inst : Academy of Sciences of USSR

Title : Luminescence Spectra of Crystals of Some Iodides.

Orig Pub : Dokl. AN SSSR, 1958, No 5, 803 - 805

Abstract : The luminescence spectra (LS) of Hg, Pb and Ag iodides consist of series of bands at the temperature of liquid air. The distances between them and their width increase together with the shift towards the long wave region. The relative intensity of the bands depends very much on the conditions, under which the sample has been prepared, and on the excitation intensity. LS of red HgI_2 consists of a narrow band ($\Delta\lambda$ about 3 mu) at about 536 mu and wide bands at about 565 and 750 mu. The most short-wave band is shifted about 3 mu from the most long-wave exciton ab-

Card : 1/2

USSR / Physical Chemistry. Crystals.

B-5

Abs Jour : Ref Zhur - Khimiya, No 8, 1957, 25922

Abstract : sorption band towards the long-wave side. LS of PbI_2 consists of a narrow band ($\Delta\lambda$ about 2 μ) at about 497 μ and wide bands at 515 and 670 μ . Stokes' shift of the narrow band is about 2 μ . There are in LS of AgI a narrow band ($\Delta\lambda$ about 2 μ) at 432 μ and wide bands at 450, 527, 650 and 780 μ . Stokes' shift is about 3 μ . A similar correspondence of absorption spectra and LS is observed also in case of CdS. The displacement of maxima of narrow radiation bands relatively to the bands of exciton absorption is nearly the half of the band width. It is concluded that the narrow radiation bands are bands of resonance luminescence and that they are caused by the collapse of the exciton state.

Card : 2/2

PLUTILL, L.

PRIKHOTKO, N.F.
24(7) 13 PHASE I BOOK EXPLOITATION SOV/1365
L'vov. Universitet

Materialy X Vsesoyuznogo soveshchaniya po spektroskopii. t. 1:
Molekulyarnaya spektroskopiya (Papers of the 10th All-Union
Conference on Spectroscopy. Vol. 1: Molecular Spectroscopy)
[L'vov] Izd-vo L'vovskogo univ-ta, 1957. 499 p. 4,000 copies
printed. (Series: Its: Fizichnyy sbirnyk, vyp. 3/8/)

Additional Sponsoring Agency: Akademiya nauk SSSR. Komissiya po
spektroskopii. Ed.: Jaxer, S.L.; Tech. Ed.: Saranyuk, T.V.;
Editorial Board: Landsterg, G.S., Academician (Resp. Ed., Deceased),
Neporent, B.S., Doctor of Physical and Mathematical Sciences,
Fabelinskiy, I.L., Doctor of Physical and Mathematical Sciences,
Kornitskiy, V.G., Candidate of Technical Sciences, Rayskiy, S.M.,
Candidate of Physical and Mathematical Sciences, Klimovskiy, L.K.,
Candidate of Physical and Mathematical Sciences, Miliyanchuk, V.S.,
A. Ye., Candidate of Physical and Mathematical Sciences.

Card 1/30

Shpol'skiy, E.V., E.A. Girdshiyanskaya, and L.A. Klimova.
Emission Spectra of Aromatic Hydrocarbons at Low
Temperatures

Gross, Ye. F., and A.A. Kaplyanskiy. Exciton Pattern
of the Spectral Curves for the Intrinsic Photoeffect
and the Exciton Luminescence Spectra in Crystals

Gross, Ye. F., B.F. Zakharchenya, and N.M. Reynov.
Zeeman Effect in the Exciton Spectrum of the
Cuprous-oxide Crystal

Pechilov, P.P. Absorption and Luminescence of Bivalent
Rare-earth Ions in Synthetic and Natural Fluorite
Crystals

Faydysh, A.M., and I. Ya. Kucherov. Migration and
Transfer of Electron-excitation Energy in Anthracene
and Naphthalene Crystals

Card 4/30

State OL
Optical Inst
in S. I. Vavilov

"APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5

APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5"

"APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5

APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R000412910009-5"

Feofilov, P.P.

48-4-18/48

SUBJECT: USSR/Luminescence

AUTHORS: Arkhangel'skaya V. A. and Feofilov P.P.

TITLE: Luminescence of Some "Pure" Salts (Lyuminestsentsiya nekotorykh "Chistykh" soley).

PERIODICAL: Izvestiya Akademii Nauk SSSR, Seriya Fizicheskaya, 1957, Vol 21, #4, p 530 (USSR)

ABSTRACT: At the liquid air temperature, luminescence spectra of some non-activated salts (HgJ_2 , PbJ_2 , AgJ , CdS) consist of a series of bands whose width increases systematically toward the longer wavelengths.

The intensity of individual bands in luminescence spectra greatly depends on the intensity of exciting light. The intensity of short wavelength bands rises with excitation intensity steeper than linear, according to the expression;

$$I \sim E^\gamma$$

where $\gamma > 1$, whereas for long wavelength bands $\gamma \leq 1$.

Card 1/2

TITLE:

48-4-18/48

Luminescence of Some "Pure" Salts (Lyuminestsentsiya nekotorykh "Chistykh" soley).

Narrow luminescence bands of Hg_2^{2+} and CdS monocrystals are fully polarized in a direction perpendicular to the crystal optical axis. Bands with longer wavelength are also polarized partially or fully in the same direction.

The luminescence duration in narrow bands does not exceed 10^{-6} sec. In bands with longer wavelengths it can attain 10^{-4} sec.

Emission spectra, as well as absorption ones, are considered to be of resonance character.

Narrow bands in emission spectra can be ascribed to decay of exciton state, although this interpretation cannot be considered as final. The report was followed by a short discussion. No references are cited.

INSTITUTION: Not indicated

PRESENTED BY:

SUBMITTED: No date indicated

AVAILABLE: At the Library of Congress.

Card 2/2

Feofilov, P.P.

48-4-40/48

SUBJECT: USSR/Luminescence

AUTHORS: Bonch-Bruyevich A. M., Tishchenko G.A. and Feofilov P.P.

TITLE: Luminescence Duration of Color Centers in Ionic Crystals
(Dlitel'nost' lyuminesstsentov tsentrov okrashivaniya v ionnykh kristallakh)

PERIODICAL: Izvestiya Akademii Nauk SSSR, Seriya Fizicheskaya, 1957,
Vol 21, #4, p 590 (USSR)

ABSTRACT: The application of a new fluorometer "GOI" possessing a high resolution capacity and relatively high sensitivity to light made it possible to investigate the luminescence duration τ of complex color centers in ionic crystals of LiF, NaF and CaF_2 . Investigated crystals were colored both photochemically, by means of hard radiation, and in the additive way (CaF_2).
In all cases the luminescence duration at room temperature was of the order of 10^{-8} to 10^{-9} sec and did not depend on the mode of coloring, intensity and wavelength of excitation light.

Card 1/2 Two types of color centers were observed in LiF crystals subjected to the action of X-rays. They differed in their